

参 考 文 献

- [1] 汤凤霞,乔长晟,宋卫东. natamycin 及其在食品中应用的研究[J]. 食品科技,2000,(6):43~44
- [2] Bas fico J C, deBas fico M Z, Chiericatti C, et al. Characterization and control of thread mould in cheese[J]. Letters in Applied Microbiology, 2001, (32): 419~423
- [3] Committee for veterinary medicinal products natamycin. The European Agency for the Evaluation of Medicinal Products Veterinary Medicines Evaluation Unit [S]. Emea / mrl/ 342/ 98-final , 1998
- [4] Stack J A, Harrison M, Perrett L L. Evaluation of a selective medium for *Brucella* isolation using natamycin [J]. Journal of Applied Microbiology, 2002, 92: 724~728
- [5] Gourama H, Bullernman L B. Effects of potassium sorbate and natamycin on growth and penicillic acid production by *Aspergillus ochraceus* [J]. Journal of Food Protection, 1998, 51(2): 139~144
- [6] Vinderrola C G, Costa G A, Regenhardt S, et al. Influence of compounds associated with fermented dairy products on the growth of lactic acid starter and probiotic bacteria[J]. International Dairy Journal, 2002(12): 579~589
- [7] 诸葛健,王正祥. 工业微生物实验技术手册[M]. 北京:轻工业出版社, 1994. 62~63, 220~222
- [8] Boer E de, Stolk-horsthuis M. Sensitivity to natamycin (pimaricin) of fungi isolated in cheese warehouses [J]. Journal of Food Protection, 1977, 40 (8): 533~536

科研简讯

绒山羊绒毛生长机理及营养调控技术研究通过鉴定

2003年12月,我校贾志海教授主持完成的“绒山羊绒毛生长机理及舍饲半舍饲营养调控技术研究”顺利通过农业部组织的专家鉴定。该项成果系统研究了绒山羊绒毛生长规律、生长机理、营养代谢及调控技术,为我国绒山羊科学化饲养提供了理论依据和技术保障。该成果达到国际先进水平,绒山羊营养调控技术达到国际领先水平。

研究成果已在内蒙古、河北、陕西等地的600万只绒山羊中推广应用,大大加快了绒山羊舍饲半舍饲进程,取得了巨大的经济效益和社会效益,绒山羊饲养区的草场退化得到了遏制,生态环境得到了恢复和改善。

(科技处供稿)